



## Corrigendum

Corrigendum to “Kinetics of the photocatalytic disinfection of *Escherichia coli* suspensions” [Appl. Catal. B: Environ. 82 (2008) 27–36]

Javier Marugán, Rafael van Grieken\*, Carlos Sordo, Cristina Cruz

Department of Chemical and Environmental Technology, ESCET, Universidad Rey Juan Carlos, C/Tulipán s/n 28933 Móstoles, Madrid, Spain


Available online 16 December 2008

The authors regret that the values of the kinetic parameters were published with errors. The reported values were obtained from the fitting of the dimensionless data  $C/C_0$ . Consequently, the values shown in Figs. 4–6, 8, 10–12 were  $k' = k/C_0$  and  $K' = K \times C_0^n$  instead of the actual values of  $k$  and  $K$ .


Therefore, the following two corrections should be considered:

- Figs. 4–6, 8, 10–12 are reproduced correctly below.
- The second paragraph in Section 3.3 should be also corrected as:

Similarly, the values of the dimensionless pseudo-adsorption constant  $K' = K \times C_0^n$  show a marked dependence on the catalysts concentration. A decrease in  $K'$  is observed as the titania concentration is increased, reaching a asymptotic minimum value of  $24.5 \pm 5.3$  for values higher than  $0.05 \text{ g L}^{-1}$  of  $\text{TiO}_2$  (Fig. 7). The variation of both parameters seems to be connected, what is confirmed by the high value of the Pearson's correlation coefficients between  $k$  and  $K$  calculated by the fitting algorithm.



**Fig. 4.** Fitting of the kinetic model to experimental data of the photocatalytic inactivation of  $10^6 \text{ CFU mL}^{-1}$  *E. coli* suspensions with different loadings of Degussa P25  $\text{TiO}_2$  (error bars calculated from eight independent bacteria counting measurements). Units of  $k$ :  $\text{CFU mL}^{-1} \text{ min}^{-1}$ ; units of  $K$ :  $\text{mL}^n \text{CFU}^{-n}$ ;  $n$  dimensionless.




**Fig. 5.** Influence of Degussa P25  $\text{TiO}_2$  loading on the kinetic constants for the photocatalytic inactivation of  $10^6 \text{ CFU mL}^{-1}$  *E. coli* suspensions (error bars calculated from two independent experiments).


DOI of original article: 10.1016/j.apcatb.2008.01.002.

\* Corresponding author. Tel.: +34 91 488 7007; fax: +34 91 488 7068.


E-mail address: [rafael.vangrieken@urjc.es](mailto:rafael.vangrieken@urjc.es) (R. van Grieken).




**Fig. 6.** Influence of Degussa P25  $\text{TiO}_2$  loading on the dimensionless pseudo-adsorption constants for the photocatalytic inactivation of  $10^6 \text{ CFU mL}^{-1}$  *E. coli* suspensions (error bars calculated from two independent experiments).




**Fig. 8.** Photocatalytic inactivation of  $10^6 \text{ CFU mL}^{-1}$  *E. coli* suspensions with two different silica-supported  $\text{TiO}_2$  photocatalysts (error bars have been omitted for clarity purposes). Lines show the fitting of the experimental data with the kinetic model represented by Eqs. (16) and (17). Units of  $k$ :  $\text{CFU mL}^{-1} \text{ min}^{-1}$ ; units of  $K$ :  $\text{mL}^n \text{ CFU}^{-n}$ ;  $n$  dimensionless.



**Fig. 10.** Influence of the bicarbonate concentration on the photocatalytic inactivation of  $10^6 \text{ CFU mL}^{-1}$  *E. coli* suspensions with  $0.05 \text{ g L}^{-1}$  of Degussa P25  $\text{TiO}_2$  (error bars have been omitted for clarity purposes). Lines show the fitting of the experimental data with the kinetic model represented by Eqs. (16) and (17). Units of  $k$ :  $\text{CFU mL}^{-1} \text{ min}^{-1}$ ; units of  $K$ :  $\text{mL}^n \text{ CFU}^{-n}$ ;  $n$  dimensionless.



**Fig. 11.** Influence of the phosphate concentration on the photocatalytic inactivation of  $10^6 \text{ CFU mL}^{-1}$  *E. coli* suspensions with  $0.05 \text{ g L}^{-1}$  of Degussa P25  $\text{TiO}_2$  (error bars have been omitted for clarity purposes). Lines show the fitting of the experimental data with the kinetic model represented by Eqs. (16) and (17). Units of  $k$ :  $\text{CFU mL}^{-1} \text{ min}^{-1}$ ; units of  $K$ :  $\text{mL}^n \text{ CFU}^{-n}$ ;  $n$  dimensionless.



**Fig. 12.** Influence of the humic acids concentration on the photocatalytic inactivation of  $10^6 \text{ CFU mL}^{-1}$  *E. coli* suspensions with  $0.05 \text{ g L}^{-1}$  of Degussa P25  $\text{TiO}_2$  (error bars have been omitted for clarity purposes). Lines show the fitting of the experimental data with the kinetic model represented by Eqs. (16) and (17). Units of  $k$ :  $\text{CFU mL}^{-1} \text{ min}^{-1}$ ; units of  $K$ :  $\text{mL}^n \text{ CFU}^{-n}$ ;  $n$  dimensionless.

## Acknowledgements

The authors thank Dr. Vitor Vilar for the discussion about the values of the kinetic parameters.